

	Individual has a positive LFT result or develops COVID-19 symptoms	Individual receives PCR test result
Index case Person who develops symptoms or has positive LFT/PCR test result	<ul style="list-style-type: none">Immediately start self-isolation and arrange COVID-19 PCR test.PCR tests can be arranged through the online portal or via 119.During self-isolation, individuals should follow 'stay at home' guidanceIf an individual does not have a PCR test within 2 days of positive LFT, they must self-isolate for 7 days minimum	<p>If PCR result is negative: end self-isolation when well and have not had a temperature for 48 hours (unless not exempt from isolation and someone else in household is symptomatic/positive)</p> <p>If PCR result is positive:</p> <ul style="list-style-type: none">Continue to self-isolate for 7 days minimum after the day of symptom onset/test (if no symptoms) – which is day 0Self-isolation can end after a lateral flow test on Day 7 if they have had 2 LFT tests taken 24 hours apart (on Day 6 and Day 7) and they are well and do not have a temperature.They should work from home if able, wear a face covering and limit contact, especially with vulnerable individuals until the end of the 10 day periodIf symptoms start whilst isolating, isolation period starts again - day of symptom onset = day 0 <p>If PCR result of index case is negative: contacts self isolating can end self-isolation if no symptoms</p> <p>If PCR result of index case is positive: action depends on age and vaccination status:</p>
Close contacts Including household contacts	<p>Action depends on age and vaccination status.</p> <p><u>Adults who are not doubly vaccinated</u></p> <ul style="list-style-type: none">Self-isolate until index case receives PCR test result. <p><u>Adult who had second vaccine dose at least 2 weeks before contact</u></p> <ul style="list-style-type: none">No requirement for self-isolation, but is strongly recommended to take an LFT test daily for 7 days prior to coming into the setting (and then continue with twice weekly LFT testing). <p><u>Child under 5.</u></p> <ul style="list-style-type: none">No requirement for self-isolation, <i>but it is a local recommendation for close contacts to be asked to PCR test and continue with regular LFT testing (if acceptable to parents).</i> PCR tests are nationally recommended for under 5s who are household contacts of positive cases. Providers may <i>wish to ask household contacts to remain away from the setting until PCR negative.</i>	<p><u>Adults who are not doubly vaccinated</u></p> <ul style="list-style-type: none">Continue to self-isolate for 10 clear days after the day of last contact with index case.If develop COVID-19 symptoms, arrange PCR test as soon as possible. <p><u>Adult who had second vaccine dose at least 2 weeks before contact</u></p> <ul style="list-style-type: none">Please see guidance for this same cohort in box to the left. <p><u>Child under 5.</u></p> <ul style="list-style-type: none">Please see guidance for this same cohort in box to the left.
	EY and childcare providers	<p><u>If the individual is in the setting when they become symptomatic:</u> Isolate individual and arrange for them to go home (or call 999 if seriously unwell). If possible, should isolate behind a closed door or 2m away from others, with an open window. Clean isolation room (and bathroom if used).</p> <p><u>Notification</u></p> <ul style="list-style-type: none">Notify cases to earlyyearsadvisors@warwickshire.gov.uk (8am to 5pm Mon - Fri) and dphadmin@warwickshire.gov.uk out of hours.LA (with UKHSA as appropriate) will complete a risk assessment, provide advice, and determine whether an Incident Management Team meeting required. <p><u>Identification of close contacts</u></p> <ul style="list-style-type: none">Providers should identify close contacts who need to self-isolate or test (see self-isolation/testing rules above and close contact definitions to right).Providers should send letter to contacts (provided by LA)

Cleaning: Please refer to detailed guidance for [cleaning of non-healthcare settings](#) and for both routine cleaning, and cleaning following an infectious person having been in your follow the key points below:

- Use disposable cloths or paper roll and disposable mop heads.
- Use, as in cleaning routine, a combined detergent/disinfectant solution at a dilution of 1,000 parts per million available chlorine (ppm av. Cl.) OR household detergent followed by disinfection (1000ppm av. Cl.) OR if an alternative combined detergent/disinfectant is used ensure it is effective against enveloped viruses and meets EN 14476 standards (including any wipes used)
- Any waste from suspected cases and cleaning of areas should be double bagged and stored for 72 hours before disposal as normal.