

Qualitative Research

Identifying strategies to increase influenza vaccination in GP practices: a positive deviance approach

Katie V Newby^{a,*}, Joanne Parsons^a, Jessica Brooks^a, Rachael Leslie^b and Nadia Inglis^b

^aCentre for Technology Enabled Health Research (CTEHR), Faculty of Health and Life Sciences, Coventry University, Coventry and ^bPublic Health Department, Warwickshire County Council, Warwick, UK.

*Correspondence to Katie V Newby, Centre for Technology Enabled Health Research (CTEHR), Faculty of Health and Life Sciences, Coventry University, Priory Street, Coventry CV1 5FB, UK; E-mail: K.newby@coventry.ac.uk

Abstract

Background. Influenza (flu) is an acute viral infection of the respiratory tract, which can lead to serious complications for individuals within at-risk groups. Evidence indicates that aspects of organization and delivery within GP practices can have an influence on the rates of flu vaccination uptake. Positive deviance is a methodological approach that facilitates identification of factors associated with high performance.

Objective. To use positive deviance to isolate factors associated with high performance by comparing GP practices achieving high and low flu vaccination uptake.

Methods. This was a qualitative study. A total of 18 practice managers and 2 GPs from 20 GP practices participated, 10 with high and low vaccination rates, respectively. Telephone interviews were conducted, audio recorded and fully transcribed. Framework Analysis was used to analyse the data.

Results. High uptake practices were more likely than low uptake practices to have a lead member of staff who demonstrated tenacity, have aspirational uptake targets, have developed and used additional prompts within their IT systems to identify eligible patients, have GPs who were opportunistically vaccinating and use phone calls as a first-line strategy to invite patients for vaccination.

Conclusions. This is the first known qualitative study to identify strategies used by UK GP practices to deliver seasonal flu vaccination programmes. It is one of few studies using the robust and novel approach of positive deviance to inform health care recommendations. This approach has offered new and more nuanced insights into GP practice factors associated high flu vaccination uptake beyond those captured through large-scale survey research.

Key words. General practice, influenza, primary health care, vaccination.

Introduction

Influenza (flu) is an acute viral infection of the respiratory tract. In healthy individuals, flu is usually self-limiting but for some it can be very serious resulting in hospitalization. In England, a free annual flu vaccination has been offered for a number of years to those aged over 65 years, those in clinical risk groups and pregnant women. In 2013, the roll-out of a programme to vaccinate all 2–16 year olds against flu was also initiated. WHO guidance

(1) indicates that developed countries should achieve a minimum of 75% uptake among the elderly population, and the European Union Council (2) recommends 75% uptake among all those at high risk. While uptake in England has been in the region of 75% for the over 65s for some years, that for clinical risk groups and pregnant women has consistently fallen well below this figure, as has uptake among the most recent cohort of children to be offered the vaccination (3).

In England, the seasonal flu vaccination programme is operated through primary care. Individual practices, typically operated by GPs, nurses and a practice manager (PM), on average vaccinate 1000 patients within a 4- to 6-week period (4). This process is managed independently with support from Public Health England (PHE) and public health colleagues working within local authorities. This is a large and complex task, requiring high levels of organization, which is performed well by international comparisons (5). In line with the Quality and Outcomes Framework (QOF) (6), GP practices receive financial incentives to achieve high uptake rates among their clinical risk groups, with pay reflecting percentage uptake beyond achieving a minimum threshold (7).

At the individual level, the decision of patients to accept an invitation for flu vaccination is influenced by a number of factors, including perception of the threat of flu and fear of side effects from vaccination (8). Evidence also indicates however that practices themselves have an influence on uptake. A study by Dexter *et al.* (4), involving the analysis of survey data from 795 practices across England, identified 7 factors associated with higher flu vaccination uptake. These included, having a lead member of staff coordinating the flu campaign, producing a written report of practice performance, having a lead member of staff responsible for identifying eligible patients, interrogating practice IT systems to identify those eligible, sending out personal invitations to patients, persisting with vaccination attempts until QOF targets are reached and the provision of flu vaccine to pregnant women by midwives. Dexter *et al.* (4) initially identified practice factors to include in the survey through interviews with staff members from six high-performing practices. The method used to identify these high-performing practices, and what constituted 'high performance', is unknown. Furthermore, the approach taken does not reliably isolate factors peculiar to high performance. For this, it is necessary to also identify the characteristics of lower-performing practices and to draw comparisons.

An emerging approach used to examine the quality of health care and make recommendations for improvement is 'positive deviance' (9). The central premise of this approach is that the knowledge about what works to improve care already exists within that community. It involves identifying and sharing practices employed by high-performing organizations and is most suitable when there is variation in performance, and where there are robust and widely utilized measures of this performance. This is the case for flu vaccination uptake, monitored centrally from returns made by individual GP practices. A methodologically strong application of positive deviance involves the identification and contrast of practices used by both high and lower performers in order to isolate factors associated with success.

There is a paucity of research reporting on the application of positive deviance to inform health care recommendations. Within the context of flu vaccination, there is one study that used this approach to identify factors associated with vaccination uptake in a clinical setting (10). This was however a US study examining how local health clinics responded during the 2009 H1N1 flu pandemic. The context of the pandemic, and the different organization of health care delivery in the USA, means that these findings have limited generalizability to the annual flu vaccination programme in the UK. This is the first known qualitative study to identify strategies used by UK GP practices to deliver seasonal flu vaccination programmes. It uses the robust and novel approach of positive deviance to isolate strategies associated with high performance by comparing GP practices achieving high and low flu vaccination uptake. This approach has

the potential to offer new and more nuanced insights into factors associated with success beyond those captured through large-scale survey research.

Methods

Institutional ethics approval was received for the study prior to commencement.

Identification of practices

The flu vaccination uptake rates (for over 65s, under 65s with chronic illness and pregnant women) for all practices in Coventry and Warwickshire were compared. Practices were sorted by percentage uptake at 31 January 2014 for each risk group. Practices with uptake above or below 75% in two or more risk groups were identified as having high or low uptake, respectively, and then ranked by overall percentage uptake, that is, combined uptake across all groups. High and low uptake practices were ranked in descending and ascending order, respectively. Finally, a small number of practices were removed where the function or population was considered to be significantly different from others, for example where primarily the practice population was homeless, as these practices were considered to be less comparable and the findings less transferable.

Recruitment

Researchers (KN and JP) contacted practices in the high and low uptake groups, working through the rank-ordered lists in descending and ascending order, respectively, and invited a senior member of staff (PM or GP) to participate. Individuals from the first 10 practices on the list in the high uptake group participated. Within the low uptake group, four practices declined to participate. Where this occurred, researchers moved on to the next practice on the list until 10 had been recruited.

Sample

Participating practices were drawn from two neighbouring local authorities with a combined area of ~2000 km² and containing a total of 139 practices. Key characteristics of the practices are presented in Table 1.

Practices were mostly located in urban areas but a small number from each group had a rural designation. Practices in both groups were situated within a range of areas of deprivation; half of high and low uptake practices were situated within the top 40% most deprived neighbourhoods; and both groups had two practices within the top 40% least deprived neighbourhoods. In both groups, there were practices with a markedly greater proportion of elderly residents or young people. Similarly, ethnic mix varied within the groups, with some practices situated within a largely White population and others within a largely non-White population. The size of practice populations varied. Practices in the high uptake group were typically smaller with a mean practice population of ~3500 (compared to 7500 in the lower group). Telephone interviews with 18 PMs and 2 GPs from 20 practices (10 high and 10 lower) were conducted.

Data collection

A semi-structured interview schedule was developed by the authors to collect information on the characteristics of each practice; types of strategies used to inform eligible patients about the flu vaccine and promote uptake; and aspects of leadership, planning and review. All interviews were conducted via telephone and audio recorded.

Table 1. Key characteristics of participating GP practices^a

Practice number	Deprivation ^b	Percentage over 65 years ^c	Percentage under 18 years ^c	Percentage non-White (British or Northern Irish) ^c	Rural and urban area classification ^d	Size of patient population (approximately)
1	13 284	25	20	3	Town and fringe	5200
2	6720	15	21	8	Urban >10k	2300
3	26 896	22	19	5	Town and fringe	3779
4	23 724	14	19	27	Urban >10k	5400
5	13 304	6	10	32	Urban >10k	4300
6	4256	9	23	50	Urban >10k	2700
7	18 099	22	17	7	Urban >10k	3200
8	3435	8	30	83	Urban >10k	2500
9	8347	7	18	54	Urban >10k	1800
10	6969	11	27	19	Urban >10k	3200
11	12 218	19	21	8	Urban >10k	4000
12	21 219	23	19	3	Village Hamlet and isolated areas	4600
13	19 669	7	25	4	Urban >10k	5600
14	864	9	31	79	Urban >10k	3600
15	10 628	19	21	31	Urban >10k	2657
16	18 395	28	9	18	Urban >10k	13 000
17	14 819	10	12	35	Urban >10k	6300
18	9514	28	13	7	Urban >10k	14 000
19	13 104	12	25	3	Town and fringe	12 000
20	22 534	13	23	6	Town and fringe	10 000

^aRows shaded in light grey represent high uptake practices, and rows shaded in dark grey represent low uptake practices.

^bBased on Index of Multiple Deprivation (IMD) sourced using Index of Deprivation 2015 Explorer. Available at <http://dclgapps.communities.gov.uk/imd/id-map.html>. IMD is given for Local Super Output Area (LSOA) in which GP practice is based. Figure represents position in ranked order of most deprived English neighbourhoods (where 1 = most deprived). Total number of total LSOAs is 32 844.

^cData based on 2011 census available at <http://www.neighbourhood.statistics.gov.uk/dissemination/LeadHome.do?m=0&s=1456823364977&cenc=1&nsjs=true&nsck=false&nssvg=false&nswid=1366>.

^dData reflect area classification for Super Output Area (SOA) in which LSOA is located. Available at <http://www.neighbourhood.statistics.gov.uk/dissemination/LeadHome.do?m=0&s=1456823364977&cenc=1&nsjs=true&nsck=false&nssvg=false&nswid=1366>.

Interviews were fully transcribed prior to analysis. At this point, each participant was given a code, for example H4 or L2, the 'H' and 'L' denoting high and low uptake practices, respectively.

Analysis

Data were analysed using Framework Analysis (11). Following a process of familiarization with the transcripts, a coding framework was developed by the first author (KN). Initially, codes reflected 'a priori' issues informed by the interview schedule; however, new codes were added as the process of indexing commenced. This involved coding each of the transcripts in turn with numerical labels, for example 2.1, 2.2, which represented each code. The framework was continuously refined as new codes emerged or as existing codes were split or merged. This resulted in six final codes. A second researcher (JB) used the framework to code a subsample of 10 transcripts to check for consistency of coding. A high level of agreement between coders was evident. Next 'charting' was undertaken. Two tables were created, one for high uptake practices and one for low uptake practices. Each table was identical, with column headings for each code and a row for each practice. A distilled summary of each participant's response in relation to each of the codes was placed within the relevant cell of the high and low tables as appropriate. Within each cell, a note indicating the page and line number of associated text was made to facilitate the identification of quotes at a later stage. This process of charting was also checked by the second researcher (JB) and once again a high level of consistency noted. Both researchers (KN and JB) then independently interpreted the data. This was done by making direct comparisons across all codes within a column, looking

for areas of difference between the high and low uptake practices. Interpretations were then compared and discussed with each other and then the wider team. The final description of themes therefore reflects a shared understanding of the data. Following analysis, recommendations for GP practices to increase their flu vaccination uptake were developed. While this was not an original aim of the study, the public health team involved felt that the learning should be shared as widely as possible. Accordingly, the recommendations were shared directly with all participating practices, and also all GP practices located within four clustered local authority areas, two of which were the authorities from which participating practices were drawn.

Results

A summary of areas of difference between GP practices in the high and low uptake groups across six finalized codes is presented below. Illustrative quotes are provided in Table 2.

Leadership

Practice managers or practice nurses were most likely to be the lead in planning and coordinating flu vaccination within practices. In the high uptake group, leadership was always the responsibility of a single individual. In the low uptake group, however, this was more fragmented, with either no clear lead in place or responsibility being shared by two or more people. While there were skilled and motivated leads within both groups, it was notable that in the high uptake group there were a few leads that were especially driven and tenacious.

Table 2. Illustrative quotes summarizing differences between high and low uptake GP practices

Code	Illustrative quote
Leadership	<p>‘So we’re so obsessed about this, or I am I guess, and I’ve been doing this job for 34 years so I can tell you that its always this big thing ... Honestly, it’s OCD [obsessive compulsive disorder]!’ (H4)</p> <p>‘She [lead] is like a dog with a bone! She really, really does want to get everybody in who’s entitled. So you know we, we do as much as we possibly, possibly can and you know we don’t really give up until it’s, we feel as if we’ve flogged it to death with some people!’ (H3)</p>
Vaccination targets	<p>‘Every year we want to do better than last year. I strongly believe that everyone in an at-risk group must be vaccinated’ (H7)</p> <p>‘As high as we possibly can, more than the previous year, that’s what we say every year’ (H9)</p>
Use of prompts on IT system	<p>‘But it’s easy to overlook that [QOF alert] because if it comes up with every single patient going in ... it’s easy for them to miss that they need the flu jab. So we just help them ... you can put a manual alert on that patient’s individual record and we’ll often do that, “Dr, needs flu jab, don’t let me keep asking xx”, you know! Or then you’ve got the [digital] appointment book and you can put messages under the name of the patient on the appointment book ... you know, with little bits of extra information like “patient is coming for”, I don’t know, “prescription review” or you might put in there instead “please can you do the flu jab while they’re in there”’ (H4)</p> <p>‘We put an alert on everybody’s record that pops up when they book an appointment, when they come and see a doctor, there’s an alert on saying “eligible for flu jab”, and then those are taken off as and when the flu jab’s given. So we continually run reports to see, at any given time, who’s entitled and hasn’t been [and had one]’ (H3)</p>
Opportunistic vaccination	<p>‘They [GPs] do [give opportunistically]; they actually have flu jabs available within their surgeries. So they will opportunistically give the flu jab to the patient’ (H5)</p> <p>‘Yes, yeah [GPs always give the vaccination opportunistically], I mean unless a patient had come in because they were having a heart attack or something like that! ... we do do a lot of opportunistic vaccinations’ (H3)</p> <p>‘You know, whilst they’re here I think people sort of like.. maybe somebody who’d say “oh I haven’t got time to book an appointment”, you know being offered it there and then I think it makes people more willing to sort of like to do it’ (H3)</p> <p>‘That’s why our doctors will do flu jabs as well, because otherwise if we just rely on the nurse doing it you might get some slip through the net’ (H6)</p> <p>‘GPs are definitely less likely themselves to give it there and then. They will normally refer it to the nurse’ (L12)</p> <p>‘They’re supposed to [laughter]... Well, most of them do, yeah, or they book them in to see the nurse’ (L19)</p>
Direct contact with patients	<p>‘We tend to adopt a calling program first, so we’ll try phoning them, and if we’ve not got an up to date phone number, in those cases, we can’t get hold of them, we will send letters’ (H6)</p> <p>‘The receptionists are very proactive. They are given lists of eligible patients and they will call them several times. We will mail out if necessary but I think the personal touch is important. Letters can be a waste of time’ (H7)</p> <p>‘We’ll run a report to identify the patients that the system is telling us should have one and haven’t had, so we’ll send a letter to all of those patients, and we generally do three letters’ (L11)</p> <p>‘That’s [sending letters] the most time efficient way of doing it because obviously there were so many people to ring you don’t have enough hours in the day to ring everybody’ (L12)</p>
Planning and promotion	<p>‘I guess there’s just, you know, there’s always a little bit more that you can do isn’t there to try, and you know we’ll certainly be positively working towards that this year to see if we can increase those numbers’ (L11)</p> <p>‘Yeah, so basically we review ... this year we wanted to sort of target the over 65s more ... I would like to improve the over 65s to get that to over 75 percent. So that will be the target this year for the nursing team. And again from that it will be; can we offer more clinics? Do we need to do weekend clinics? It’s how we actually all sit down and discuss that in July’ (L12)</p> <p>‘No, no [we don’t review], because, I don’t know if you saw last year’s figures, but we’re usually one of the highest, usually first or second in the city and that’s been for quite a number of years now, so we’re happy with what we’re doing’ (H9)</p> <p>‘If it works why change it? If we were struggling to hit our targets we’d do other things’ (H10)</p>

Vaccination targets

Most GP practices had flu vaccination targets and largely these were in line with the QOF minimum thresholds for clinical risk groups, and to achieve 75% uptake among the over 65s. There were several practices, however, within the high uptake group that had more aspirational targets that appeared to be more public health focussed than financially driven.

Use of prompts on IT system

Low uptake practices tended to use standard prompts built into their IT systems to alert the user that a patient was eligible for a flu vaccination. High uptake practices on the other hand made more sophisticated use of their systems by hand entering additional prompts.

Opportunistic vaccination

There was a clear distinction here between groups. In high uptake practices, GPs were reported to deliver the vaccination opportunistically, not allowing opportunities to vaccinate pass. There was an appreciation that this approach could capture some of the more ambivalent patients who may not otherwise book a separate appointment. There were a number of comments from those in the low uptake group, however, which indicated that this was not the culture within their organization.

Direct contact with patients

At most practices, all eligible patients had at least one form of direct contact about the flu vaccination. Within the high uptake practices, initial contact was often made using several methods simultaneously,

one of which was likely to be by telephone. In low uptake practices, however, first contact was usually by a single method. Telephone calls were also sometimes used, but these were more likely to be used to 'mop up' the remaining patients.

Planning and promotion

It was clear that the flu vaccination programme placed a strain on GP practices and that high levels of planning and organization were required to manage effective delivery. Most practices reported having a review and planning meeting sometime between June and August thus requiring that any input from supporting organizations is provided at or before this time. It was clear that in the low uptake group, a number of practices wished to identify where they could make improvements and also to develop and test out new strategies. In the high uptake group on the other hand, there were a few practices that reported that they were not exploring other strategies, despite vaccination uptake still being low in some at-risk groups, particularly for pregnant women, and 2–3 year olds.

Discussion

Using the approach of positive deviance, we identified GP practices with diametrically opposed flu vaccination uptake rates and compared their in-house approaches and strategies to the vaccination of their patients. Five clear and distinct factors emerged. High uptake practices were more likely than low uptake practices to have a single lead member of staff who demonstrated tenacity, to have aspirational uptake targets, to have developed and used additional prompts within their IT systems to identify eligible patients, to have GPs who were opportunistically vaccinating and to use a combination of methods including phone calls as a first-line strategy to contact and invite patients for their vaccination.

The findings of this study should be considered in light of its strengths and weaknesses. The approach used to identify practice characteristics and strategies associated with high flu vaccination uptake is robust and ensures that factors common across the board are not erroneously selected. The method used to identify high and low uptake practices was also strong, ensuring that practices had relatively high flu vaccination uptake for all at-risk groups, rather than just a high overall uptake figure. Practices were also drawn from a reasonably large geographical area, with diverse neighbourhoods including a mix of rural and urban locations, and both high and low levels of deprivation. A weakness of the study was ceasing data collection on pragmatic grounds rather than on the achievement of saturation. It is possible that including further practices would have enabled further differences between high- and lower-performing groups to emerge. A further weakness was the absence of information on the number of whole time equivalent (WTE) GPs, nurses and other supporting staff within each practice. This could have been used along with data on the size of each practice population to provide an indication of the level of patient demand. This is important as patient demand may have affected each team's ability to perform strategies to increase vaccination uptake.

The findings from this study share similarities to those by Dexter *et al.* (4), which have been used to develop a checklist of recommend strategies for GP practices to employ in the annual flu plan produced by Public Health England (see ref. (12) for the most recent edition). There are however some additional and more nuanced factors that warrant attention. Dexter *et al.* (4) identified that having a lead member of staff arranging the vaccination campaign was associated with increased uptake. In the present study, all of the high uptake

practices had a single lead member of staff. In the low uptake practices, however, leadership was somewhat fragmented with either no lead or the responsibility shared across members of the team. What also set the high uptake practices apart was having a lead who was particularly driven and determined to achieve their uptake target. Leadership in the high uptake practices pertained not only to key functions such as ordering in sufficient supply of vaccine and managing their cold storage, as were common among their counterparts, but also to the setting of targets, running reports to identify eligible patients, coordinating and enthusing staff to work together and regularly reviewing and reporting on progress.

In the high uptake practices, an aspirational uptake target was more likely to be reported than in the low uptake practices. The target for low uptake practices was usually the achievement of QOF benchmarks. High uptake practices indicated that there was a continual push to improve performance and to do better for their patients. Dexter *et al.* (4) asked respondents to indicate at what point they ceased vaccinating each year. Continuing vaccination until QOF targets had been reached rather than stopping sooner, such as when vaccine supply ran out, was associated with greater uptake. They concluded that the pursuit of QOF targets may be a driving force behind higher uptake levels. In the present study, those leading vaccination within GP practices were almost without exception PMs or practice nurses who are unlikely to personally benefit from any related financial reward. While QOF targets may be working to motivate performance up to a threshold value, it appears that more aspirational attainment is driven by motivations beyond financial gain.

This study corroborates the finding of Dexter *et al.* (4) that higher uptake is associated with sophisticated use of IT systems and the addition of in-house prompts to alert practice staff to patient eligibility for flu vaccination. Exceptional examples were uncovered of lead members of staff interrogating their systems on a daily basis, identifying patients due to visit and then coordinating the response of staff to ensure the opportunity for vaccination was not lost. Dexter *et al.* (4) found higher flu vaccination uptake to be associated with direct patient contact. In the present study, a variety of methods was used across the two groups to contact patients. What set the high uptake practices apart was their use of multiple methods, including making telephone calls, as a first-line strategy. Telephoning patients may be particularly effective due to the personal nature of the invitation, because it reduces burden on the patient and because it may capture the more ambivalent patients who may not otherwise be sufficiently motivated to book an appointment. Of note, the low uptake practices tended to be larger in size than the high uptake practices, which may make this approach less feasible.

A factor that was isolated as unique to high uptake practices in the present study was the tendency for GPs to opportunistically vaccinate. While this was reported by high and low uptake practices, those in the former group were doing it more consistently and there was the suggestion that they recognized the importance of this in order to avoid some patients, particularly, the more ambivalent ones, 'slipping through the net'. Opportunistic vaccination by GPs was not investigated by Dexter *et al.* (4). Our findings suggest that this may be an additional distinguishing factor associated with higher uptake.

Conclusions

The above findings lead to primary understandings that are of value to GP practices and public health officials tasked with monitoring and improving flu vaccination uptake. Strategies, including

providing in-house training on how to exploit the available functions within practice IT systems, training and supporting administrative staff to invite patients for vaccination by telephone and developing a culture and systems that support GPs to embed vaccination into routine appointments, all have the potential to positively impact upon vaccination rates. Furthermore, professional development and training of primary care staff that encourages individuals to lift expectations above routine targets and achievement would be valuable.

The findings of the present study have been contrasted with those of Dexter *et al.* (4) to demonstrate the complementarity of approaches. On the whole, the factors associated with success across the two studies overlap; however, the approach adopted by the present study more precisely defines the criteria. Large-scale survey studies have an important place, enabling the strength of association between practice factors and vaccination uptake to be statistically derived and compared. However, it should be recognized that in doing so these factors are potentially reduced to proxy measures. Breaking down the aspects of success that are salient, and in particular isolating those that are unique to high-performing organizations, in addition to larger-scale quantitative studies, is a methodologically sound and rigorous approach. This combination of methods enriches understanding and provides solid ground on which to inform policy change.

Acknowledgements

The authors would like to sincerely thank each participant for their time and willingness to provide information on flu vaccination within their practices. The positive deviance approach only works when organizations are open to sharing and learning as demonstrated by all of those involved.

Declaration

Funding: Warwickshire County Council and Coventry City Council.
Ethical approval: received institutional ethical approval prior to commencement.
Conflict of interest: none.

References

1. World Health Organisation. *Prevention and Control of Influenza Pandemics and Annual Epidemics*. Geneva, Switzerland: WHO, 2003. http://www.who.int/immunization/sage/1_WHA56_19_Prevention_and_control_of_influenza_pandemics.pdf (accessed 29 February 2016).
2. Council of the European Union. *Council recommendation of 22 December 2009 on seasonal influenza vaccination (2009/1019/EU)*. OJEC, 2009, pp. 348(71e2). http://www.epha.org/IMG/pdf/Council_Recommendation_on_seasonal_flu_vaccine.pdf (accessed 29 February 2016).
3. Public Health England. *Surveillance of Influenza and Other Respiratory Viruses in the United Kingdom: Winter 2014 to 2015*. 2015. https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/429617/Annualreport_March2015_ver4.pdf (accessed 29 February 2016).
4. Dexter LJ, Teare MD, Dexter M, Siriwardena AN, Read RC. Strategies to increase influenza vaccination rates: outcomes of a nationwide cross-sectional survey of UK general practice. *BMJ Open* 2012; 2: e000851.
5. Blank PR, Schwenkglenks M, Szucs TD. Vaccination coverage rates in eleven European countries during two consecutive influenza seasons. *J Infect* 2009; 58: 446–58.
6. Gillam S, Siriwardena A. *The Quality and Outcomes Framework: QOF Transforming General Practice*. Oxford, UK: Radcliffe Publishing Ltd, 2010.
7. NHS Employers. 2015/16 General Medical Services (GMS) Contract Quality and Outcomes Framework (QOF): Guidance for GMS Contract 2015/16. 2015, No. 03099. <http://www.nhsemployers.org/~media/Employers/Documents/Primary%20care%20contracts/QOF/2015%20-%202016/2015%2016%20QOF%20guidance%20for%20stakeholders.pdf> (accessed 29 February 2016).
8. Bish A, Yardley L, Nicoll A, Michie S. Factors associated with uptake of vaccination against pandemic influenza: a systematic review. *Vaccine* 2011; 29: 6472–84.
9. Bradley EH, Curry LA, Ramanadhan S *et al.* Research in action: using positive deviance to improve quality of health care. *Implement Sci* 2009; 4: 25.
10. Klaiman T, O'connell K, Stoto M. Local health department public vaccination clinic success during 2009 pH1N1. *J Public Health Manag Pract* 2013; 19: E20–6.
11. Richie J, Spencer L. Qualitative data analysis for applied policy research. In: Huberman M, Miles MB (eds). *The Qualitative Researcher's Companion*. California: Sage Publications, 2002, pp. 173–94.
12. Public Health England. *Flu Plan: Winter 2015/16*. 2015. https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/418038/Flu_Plan_Winter_2015_to_2016.pdf (accessed 29 February 2016).